

Insulating gap in FeO: correlations and covalency

I. I. Mazin^{a,b} and V.I. Anisimov^{a,c}

^a*Max-Planck-Institut für Festkörperforschung, Stuttgart D-70569, Germany*

^b*Geophysical Laboratory and Center for High-Pressure Research, Carnegie Institution of Washington, 5251 Broad Branch Rd. N.W., Washington, D.C. 20005-1305 U.S.A.*

^c*Institute for Metal Physics, Ekaterinburg, Russia*

We report calculations of the electronic structure of FeO in the LDA and LDA+U approximation with and without rhombohedral distortion. In both cases LDA renders an antiferromagnetic metal, and LDA+U opens a Hubbard gap. However, the character of the gap is qualitatively different in the two structure, and the difference can be traced down to underlying LDA band structure. An analysis of the calculations gives a new insight on the origin of the insulating gap in 3d monoxides and on the role of the $\mathbf{k}$ -dependency of U, missing in the contemporary LDA+U method.

It is a well-known fact that conventional band structure calculations incorrectly give metallic ground state for the intermediate 3d transition metal monoxides, CoO and FeO. Before the high- T_c cuprates entered the scene, this had been often considered as the most notable failure of the Local Density Approximation (LDA). In the last decade a number of extensions of the density-functional theory were suggested, which, in different manners, led to insulating ground states for this compounds. Most successful were various flavors of the Self-Interaction Corrected (SIC) LDA¹, LDA+U², and the Orbital Polarization Correction in a crystal field basis³. Interestingly, apart from the gap itself, LDA appears to do as good, and sometimes a better job, than these sophisticated extensions, especially when Generalized Gradient Correction to the conventional LDA is taken into account. Structural properties are reproduced very well, including rhombohedral distortion in FeO⁴, and its increase with pressure⁵. Moreover, angular resolved photoemission renders the bands more similar to the LDA bands, than to those in other calculational scheme (except for the narrow range near the gap)⁷. The magnitude of the magnetic moment, which is often believed to be the first indication of an LDA failure, is nearly exact for the spin moment in CoO ($2.4 \mu_B$) and in FeO ($3.5 \mu_B$, assuming no orbital moment). On the contrary, the gap-improving calculation tend to overestimate the Fe moment, especially when the orbital moment is included (i.e., there is a tendency³ to underestimated the crystal field quenching of the orbital moment). It seems, then, that the main problem in the LDA is purely spectroscopical (quite in the spirit of the density functional theory), namely non-existence of the gap. It is worth noting that an important common feature of the LDA and the corrected schemes mentioned above is substantial width of the metal d -bands (for instance, in LDA the width of Fe t_{2g} band at normal pressure is about 1.4 eV). This should be contrasted with the popular analysis of the electronic structure of 3d-oxides in terms of separate levels of a width less than Jan-Teller energy and spin-orbital coupling⁶.

The fact that LDA gaps are always too small is well understood. Mathematically it appears as the density-derivative discontinuity of the Kohn-Sham potential in the DFT. The physics of this discontinuity may be different, but for transition metal oxides (NiO, MnO) it is usually associated with the Mott-Hubbard repulsion³. In view of this, it has always been much more disturbing to have wrongly a metallic behavior in LDA calculation, than just to have a wrong gap. As formulated by Norman³, “one would like to obtain a gap at the level of a density functional calculation (no matter how small) so as to define the Mott-Hubbard correction in an unambiguous fashion”. Moreover, even in a case when LDA does not give a gap, but gives reasonable band structure except for the immediate vicinity of the Fermi level, and correctly describes delicate features of the ground state, like magnetoelastic interactions, it is desirable to have a correction scheme which does not destroy the LDA bands completely, but rather corrects them in a systematic manner. Unfortunately none of the schemes above acts in such a way. In this paper we shall analyze the results of the rotationally invariant LDA+U⁹ calculations for FeO in more details than is usually done, and compare them with the standard LDA calculations, paying particular attention to the process of the LDA+U gap opening in cubic and rhombohedral structure. We will see that the ground state in the LDA+U approach is intimately related with the underlying LDA band structure, although LDA+U cannot fully account for the bands hybridization effects, which seem to be quite important here. We will argue that none of the existing “corrected LDA” schemes (nor the straight LDA) correctly describes the insulating ground state in FeO and similar compounds. On the other hand, non-local schemes similar to the GW approximation may be able to provide a qualitatively correct description. It is worth noting that none of the existing “extended-LDA” calculations has taken into account such an important factor as the distortion from the ideal cubic NaCl-type structure, which is associated in 3d-monoxides with the onset of magnetic ordering.

Electronic structure of an isolated Fe^{++} ion in a cubic field is described in the high-spin state by the following scheme⁸: The spin-up d -states are all filled, and separated from the spin-down states by the exchange splitting E_{ex} . The partially occupied spin-down states are split by the crystal field, so that $\epsilon(e_g) - \epsilon(t_{2g}) = \Delta \ll E_{ex}$. There is one electron in the t_{2g_l} state, which is triply degenerate. Antiferromagnetic FeO has close-packed (111) planes of Fe ions with the same spin, which alternate with the similar planes with the opposite spin, thus lowering the symmetry to rhombohedral one. In the rhombohedral field, there is one linear combination of the t_{2g} states, namely $A_{1g} = (xy + yz + zx)/\sqrt{3}$, which can also be written as $3Z^2 - r^2$, where Z is parallel to $[111]$ (here and below we shall use lower case symbols for the states classification in cubic symmetry and upper case symbols for the rhombohedral nomenclature). The four other states have the same symmetry E_g (in the rhombohedral nomenclature), but when the deviation from the cubic symmetry is small one can speak about the upper two levels, E_g'' , and the lower two levels, E_g' , which are close to the A_{1g} also originate from the cubic t_{2g} states, and are separated from E_g'' by approximately Δ . In the LDA calculation, when the crystal symmetry is still cubic, although the magnetic ordering is rhombohedral, the splitting between E_g' and A_{1g} is small, much smaller than their bandwidths, so that they merge into one band (corresponding to the cubic t_{2g} band), which is necessarily metallic. Note that although the magnetism in this system appears due to the indirect exchange and is determined by the O-Fe pd interaction, the width of the t_{2g} band is mainly due to the Fe-Fe $dd\sigma$ overlap. Fe-O hopping for this band is mainly $pd\pi$ and weak. The situation in CoO with two t_{2g} electrons is very similar.

There were several successful attempts to obtain an insulating state in FeO^{1-3} . Interestingly, all these approaches give insulating gap in a fair agreement with the experiment, but all because of different reasons. SIC calculations¹ favor d -bands in pure orbital states, that is, undermining the role of crystal field and quenching of orbital moment. The effect on the occupied d -states is thus extremely strong (of the order of 1 Ry) and all four oxides come out as pure charge transfer insulators. This is in contradiction with the general experimental indication that the character of the band gap changes from predominantly Mott-Hubbard to predominantly charge-transfer when going from MnO to NiO. Large orbital moment is obtained for FeO, in contradiction with the experiment (an argument is usually made that the experimental number may be incorrect because poor sample quality). It is also worth noting that the SIC formalism was initially invented as a remedy of the LDA in the direction of the exact, self-interaction free density functional. It is hard to imagine, however, that the exact density functional theory with its orbital-independent one-electron potential can unquench the orbital moment.

The orbital-corrected functional used by Norman³ has

a similar problem. The correct many-body solution for an isolated ion with unfilled t_{2g} shell in a cubic field⁸ has an energy contribution proportional to the total angular momentum. This term, however, does not favor a specific direction of the momentum, that is, does not include the projections. The only interaction which does unquench the orbital momentum is spin-orbit, and it is relatively weak. Norman's correction has the same functional form as the exact quantum chemical expression⁸, but substitutes the total momentum by its projection on the quantization axis. This orbital moment projection dependent term acts in a way similar to the spin-orbit coupling, but with a much larger magnitude (of the order of exchange splitting J). Thus in FeO the occupied spin-down d -band is too close to a pure $m = 1$ state, that is, $xz + iyz$.




FIG. 1. (a) LDA band structure without a rhombohedral distortion. The bars show the relative A_{1g} character of the corresponding states; (b) LDA+U bands for the same crystal structure. Note the bottom of the Fe(s) band at the Γ point.

The corresponding band structure is shown on Fig.1, where the LDA+U, equally successful in predicting a band gap, forms a completely different insulating state. There is no mechanism for unquenching the orbital moment, apart from the spin-orbit coupling, neglected in Ref.² and other LDA+U calculations. LDA+U starts from the straight LDA bands, where the t_{1g} bands are

strongly mixed, the A_{1g} (in rhombohedral notations) orbital being slightly more occupied than each of the two E'_g orbitals. When formulated in rotationally invariant form⁹, LDA+U tends to apply the positive U correction to less occupied E'_g orbitals, making them less and less filled (potential U correction for more filled A_{1g} -orbital is negative), and eventually splits off the A_{1g} band, forming a gap between it and the E'_g bands. Sufficiently large U (we used the empirical value of $U=5.1$ eV¹⁰, which gives a good value for the gap; constrain LDA calculations¹¹ yield a somewhat larger $U=6.8$ eV) pushes the occupied A_{1g} band down close to, and for $U=6.8$ eV right into, the $O(p)$ -bands manifold, and the unoccupied E'_g bands up above the bottom of the Fe(s) band. This seems to be in qualitative agreement with the photoemission experiments, which show the top of the valence band to be of the mixed $O2p$ -Fe $3d$ character¹², and with the optical experiments, which indicate a weak absorption between 0.5 and 2.0 eV, assigned to the (pd) – s transitions, and strong absorption edge at 2.4 eV due to transitions into the Fe(d) band¹³. In our calculations, the minimal gap opens between the $O(p)$ –Fe(d) band and the Fe(s) band at 1.3 eV and transitions into the Fe(d) band start at 2.2 eV.

LDA and the LDA+U bands refer to the cubic FeO (here and below we show results of the ASA LMTO¹⁵ calculations with the unit cell volume 256 bohr³/FeO, close to the LDA equilibrium volume⁵). Since only 1 orbital, A_{1g} , is occupied (see Table II), there is no orbital moment. One can expect that including weak (a few mRy) spin-orbit coupling will create a small orbital moment, but hardly one comparable with the spin magnetization. It is interesting that the way the gap opens in LDA+U in the cubic structure is very “LDA-like”: The only way to open a gap in an effective one-electron approximation is to split the t_{2g} band by (magnetic, in this case) rhombohedral symmetry and to occupy the A_{1g} orbital for FeO and the two E'_g orbital in CoO. In fact, this is exactly what happened in the LDA calculations of Dufek et al¹⁴, who used an exotic LDA functional, which gave poor total energies but did open gaps in both compounds.

Overall this seems to be a physically satisfactory description of the gap opening. However, this is not as straightforward as one may think. To show the problem, let us compare the calculations in the cubic structure with those in the rhombohedrally distorted structure (Fig. 2; we used the LDA equilibrium distortion, calculated in Refs. 5; experimental distortion is smaller, but increases with the pressure). To understand the result, one should keep in mind that dispersion of the t_{2g} bands is mainly due to direct $dd\sigma$ hopping between the like spins, which can be easily verified in the LMTO-TB method by removing oxygen orbitals from the basis set, or by looking directly on the corresponding elements of the LMTO-TB Hamiltonian. We observe that the distortion increases the t_{2g} bandwidth (due to decreased distance between the like-spin ions), but mainly at the

expense of the E'_g -like bands. The dispersion of the A_{1g} band decreases instead, thus leading to a decreased occupancy of the A_{1g} -like state (because the whole t_{2g} manifold is less than half filled). One can see this by comparing the occupancy matrices in the cubic (Table I) and in the distorted (Table III) structure on the first iteration, *i. e.*, before the effect of U . While in the cubic structure the A_{1g} state was the mostly occupied one (with the two E'_g states close next), in the distorted structure the E'_g states are twice more occupied than the A_{1g} one.




FIG. 2. The same as Fig.1, but with the rhombohedral distortion of 6%.

The LDA+U code we used is rotationally invariant⁹, which means that the program can apply the U corrections to the states which are arbitrary linear combinations of the d -orbitals with various m . The coefficients of this combination are chosen self-consistently so that they optimize the total energy including Coulomb one. In the case of the rhombohedrally distorted FeO it is energetically unfavorable to occupy the A_{1g} orbital, but not because of the Hubbard energy, but because of the one-electron energy: the same reason why LDA deoccupied this orbital after distortion. How does LDA+U handles this situation during the self-consistency cycle? It is left with the only possibility to make a linear combination of the E'_g orbitals with a possible admixture of the E''_g states, as much as the crystal field allows that, to make

it a separate band and to open a gap between this band and the rest of the spin-minority d -bands (Table IV).

One may feel it unphysical that the method converges to qualitatively different ground states with and without distortion. This fact suggests that probably the correct solution should be intermediate between the two LDA+U stationary points, a completely full and a completely empty A_{1g} state. Indeed, a closer look at the LDA band structure of the Fig.2a shows that along the whole Γ -Z (111) direction the A_{1g} band lies entirely below the Fermi level. In the other parts of the Brillouin zone (points L and F) it is completely above the Fermi level. Thus, a U matrix acting on an unoccupied A_{1g} orbital has to overcome the natural LDA tendency along the Γ -Z line, although it is in accord with the LDA tendencies in the rest of the zone.

From this consideration one can see that the problem with the LDA+U is that the U -matrix is $\mathbf{k}$ -independent: it is a matrix in the orbital space, $U_{mm'}$, $-2 \leq m \leq 2$, and this matrix does not know whether the particular d -electron is in a bonding or antibonding interaction with the neighboring sites. While this may be a good approximation for extremely localized electrons, in such materials as FeO one cannot neglect the fact that d -orbitals extend into neighboring sites. In the LMTO scheme the tails of each orbital, penetrating into an atomic sphere of a neighboring site, is re-expanded and, depending on the LMTO flavor (“representation”), may or may not appear in the density matrix at the same site as its head¹⁶. In the conventional LDA+U scheme, as well as in all other correction schemes discussed above, only the heads of the LMT orbitals are subjects to a correction. On the other hand, a more realistic correction would also apply to that part of a d -orbital that penetrates into the neighboring spheres. In other words, a physically correct scheme for implemented such a correction should be essentially non-local. Hartree-Fock like schemes, similar to the GW approximation¹⁷, should be able to reproduce the correct physics. For instance, one can expect that the occupied band that appears to be purely A_{1g} in the cubic case and purely E'_g in the distorted case, would have a mixed character, being more A_{1g} -like close to the Γ -Z line. Correspondingly, the effective U -correction matrix would be different when applied at different $\mathbf{k}$ -points. Another interesting alternative is offered by non-Kohn-Sham versions of the Density Functional Theory, where a non-local (screened Hartree-Fock like) contribution to the total energy is singled out together with the non-interacting kinetic energy, and the rest is treated in an LDA. Such schemes retain the good accuracy of the total energy calculations in the LDA, and improve the excitation energies as well. It is worth noting that underestimation of the hybridization tendencies inside the t_{1g} band in LDA+U and other schemes is likely to lead to a wrong energetics and spoil the agreement of structural properties, calculated in LDA⁵, with the experiment. More detailed discussion of structural and magnetoelastic properties of FeO will be published elsewhere¹⁸.

To summarize, we report conventional LDA calculations and rotationally-invariant LDA+U calculations for antiferromagnetic FeO, both in the cubic and in a rhombohedrally distorted structure. In both cases LDA+U opens gaps, but these gaps are of completely different character in the two structures, and also different from the gaps appearing in other “corrected-LDA” schemes, which, in turn, differ by their physical nature from each other. We believe that this is a consequence of the local character of the Mott-Hubbard correction in the conventional LDA+U method, and neither straight LDA, nor “corrected-LDA” methods provide a proper physical description of the gap formation. We argue that the ultimate method must take into account the non-local character of the Coulomb repulsion.

We are thankful to O.K. Andersen, R.E. Cohen, D.I. Khomskii, A.I. Liechtenstein, and W.E. Pickett for useful discussion. We acknowledge the hospitality of the Max-Planck-Institut für Festkörperforschung, where most of this work was performed. One of us (VIA) was in part supported by Russian Foundation for Fundamental Research (grant RFFI 96-02-16167).

-
- ¹ A. Svane and O. Gunnarsson, Phys. Rev. Lett. **64**, 1162 (1990); Z. Szotek, W.M. Temmerman, and H. Winter, Phys. Rev. B **64**, 4029 (1993).
 - ² V.I. Anisimov, J. Zaanen and O.K. Andersen, Phys. Rev. B **44**, 943 (1991);
 - ³ M.R. Norman, Phys. Rev. B **44**, 1364 (1991); Int. J. Quant. Chem. Symp. **25**, 431 (1991).
 - ⁴ CoO has tetragonal distortion, but no full-potential LDA calculations for tetragonal phase were reported in the literature.
 - ⁵ D.G. Isaak, R.E. Cohen, M.J. Mehl, and D.J. Singh, Phys. Rev. B **47**, 7720 (1993); D.M. Sherman and H.J.F. Jansen, Geophys. Lett. **22**, 1001 (1995).
 - ⁶ J.B. Goodenough, Phys. Rev. B **171**, 465 (1966); K.I. Kugel and D.I. Khomskii, Sov. Phys. Usp., **25**, 231 (1982).
 - ⁷ Z.X. Shen *et al*, Phys. Rev. Lett. **64**, 2442 (1990); Phys. Rev. B **42**, 1817 (1990); *ibid*, B **44**, 3604 (1991).
 - ⁸ J.S.Griffith, *The theory of transition metal ions*, Cambridge, 1971
 - ⁹ A.I. Lichtenstein, J. Zaanen, V.I. Anisimov, Phys. Rev. B **52**, R5467 (1995)
 - ¹⁰ J. Zaanen and G.A. Sawatzky, J. Solid State Chem. **88**, 8(1990)
 - ¹¹ V. I. Anisimov and O. Gunnarson, Phys. Rev. B **43**, 7570 (1991).
 - ¹² P.S. Bagus, C.R. Brundle, T.J. Chuang, K. Wandelt, Phys. Rev. Lett. **39**,1229 (1977)
 - ¹³ I. Balberg, H.L. Pinch, J. Magn. Magn. Mater. **7**, 12 (1978)
 - ¹⁴ P. Dufek, P. Blaha, and K. Schwarz, Phys. Rev. B **50**, 7279 (1994);
 - ¹⁵ We used the LMTO-TB-ASA code *Stuttgart 4.7*.

- ¹⁶ O.K. Andersen and O. Jepsen, Phys. Rev. Lett., **53**, 2571 (1984).
¹⁷ F. Aryasetiawan and O. Gunnarsson, Phys. Rev. Lett., **74**, 3221 (1995).
¹⁸ I.I. Mazin, R.E. Cohen, and D.G. Isaak, to be published.

TABLE I. Orbitals and their occupations without rhombohedral distortion, for $U = 0$. Coordinates system corresponds to the rhombohedral symmetry: z-axis is perpendicular to the ferromagnetic Fe planes, y-axis points towards the nearest neighbor Fe with the like spin. The character of the orbitals is, in the order of the table, $2 \times E'_g$, $2 \times E''_g$, A_{1g}

occupation	xy	yz	$3z^2 - 1$	xz	$x^2 - y^2$
.17	-.03	.03	.00	-.65	.76
.17	.75	-.66	.00	-.03	.03
.31	.60	.69	.00	.30	.26
.33	-.26	-.30	.00	.69	.60
.39	.00	.00	1.00	.00	.00

TABLE II. Orbitals and their occupations without rhombohedral distortion, for $U = 5.1$. The character of the orbitals is $2 \times E''_g$, $2 \times E'_g$, A_{1g}

occupation	xy	yz	$3z^2 - 1$	xz	$x^2 - y^2$
.04	-.10	-.10	.00	.68	.72
.04	.72	.68	.00	.10	.10
.10	.53	-.57	.00	-.46	.43
.10	-.43	.46	.00	-.57	.53
.90	.00	.00	1.00	.00	.00

TABLE III. Orbitals and their occupations with 6% rhombohedral distortion, for $U = 0$. The characters are $2 \times E'_g$, A_{1g} , $2 \times E''_g$

occupation	xy	yz	$3z^2 - 1$	xz	$x^2 - y^2$
0.17	-.55	.84	.00	-.02	.01
0.17	-.01	.02	.00	.83	-.56
0.19	.00	.00	1.0	.00	.00
0.38	.84	.55	.00	.00	-.00
0.39	-.01	-.01	.01	.56	.83

TABLE IV. Orbitals and their occupations with 6% rhombohedral distortion, for $U = 5.1$. The characters are A_{1g} , E''_g , $2 \times E'_g$, A_{1g} , E''_g

occupation	xy	yz	$3z^2 - 1$	xz	$x^2 - y^2$
.04	-.02	.00	.99	.14	.04
.04	.96	.28	.02	.01	.01
.10	-.28	.96	.00	.02	-.01
.10	.01	-.02	-.09	.83	-.56
.86	-.01	.00	-.12	.55	.83